

N,N'-Bis(2-aminobenzyl)ethane-1,2-diaminium dinitrate

Luis Ángel Garza Rodríguez,^a Perla Elizondo Martínez,^b Sylvain Bernès,^{b*} Blanca Nájera Martínez^b and Nancy Pérez Rodríguez^b

^aUniversidad Regiomontana A.C., 15 de Mayo 567 Pte., Monterrey, Nuevo León CP 64000, Mexico, and ^bUniversidad Autónoma de Nuevo León, UANL, Facultad de Ciencias Químicas, Av. Universidad S/N, Ciudad Universitaria, San Nicolás de los Garza, Nuevo León CP 66451, Mexico

Correspondence e-mail: sylvain_bernes@Hotmail.com


Received 13 September 2013; accepted 7 October 2013

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.012\text{ \AA}$; R factor = 0.061; wR factor = 0.180; data-to-parameter ratio = 8.9.

In the title salt, $\text{C}_{16}\text{H}_{24}\text{N}_4^{2+} \cdot 2\text{NO}_3^-$, both the cation and anion are placed in general positions, although the cation displays non-crystallographic inversion symmetry, with the aliphatic chain extended in an all-*trans* conformation. The benzene rings are almost parallel, with a dihedral angle between their mean planes of $3.3(6)^\circ$. The nitrate ions are placed in the vicinity of the protonated amine groups, forming efficient $\text{N}-\text{H}\cdots\text{O}$ inter-ion hydrogen bonds. Each nitrate ion in the asymmetric unit bridges two symmetry-related cations, forming an $R_4^4(18)$ ring, a common motif in organic ammonium nitrate salts. This results in the formation of chains along [010] with alternating cations and anions. The neutral amine groups are involved in slightly weaker $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds with the nitrate O atoms, and there are also a number of $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds present. The resulting supramolecular structure is based on a two-dimensional network extending in the *ab* plane.

Related literature

For the structure of the free neutral amine, see: Rodríguez de Barbarín *et al.* (2007). For the *p*-toluenesulfonate salt of the title cation, see: Garza Rodríguez *et al.* (2011). For related diammonium nitrate salts featuring $R_4^4(18)$ motifs, see: Liu *et al.* (2007); Yang *et al.* (2007). For supramolecular motifs nomenclature, see: Etter (1990). For the synthesis of the title salt, see: Garza Rodríguez (2010).



Experimental

Crystal data

$\text{C}_{16}\text{H}_{24}\text{N}_4^{2+} \cdot 2\text{NO}_3^-$	$V = 1912.1(18)\text{ \AA}^3$
$M_r = 396.41$	$Z = 4$
Orthorhombic, Pna_2	Mo $K\alpha$ radiation
$a = 11.041(5)\text{ \AA}$	$\mu = 0.11\text{ mm}^{-1}$
$b = 5.760(4)\text{ \AA}$	$T = 298\text{ K}$
$c = 30.069(13)\text{ \AA}$	$0.60 \times 0.20 \times 0.20\text{ mm}$

Data collection

Siemens P4 diffractometer	$R_{\text{int}} = 0.053$
4371 measured reflections	3 standard reflections every 97
2473 independent reflections	reflections
1501 reflections with $I > 2\sigma(I)$	intensity decay: 1.5%

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.061$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.180$	$\Delta\rho_{\text{max}} = 0.34\text{ e \AA}^{-3}$
$S = 1.61$	$\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$
2473 reflections	
278 parameters	
13 restraints	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N9—H9A \cdots O23 ⁱ	0.94 (4)	1.87 (3)	2.794 (8)	167 (7)
N9—H9B \cdots O24	0.95 (5)	1.96 (5)	2.879 (7)	164 (5)
N12—H12A \cdots O27	0.92 (3)	1.88 (3)	2.787 (8)	168 (8)
N12—H12B \cdots O28 ⁱ	0.91 (5)	1.95 (5)	2.862 (8)	176 (9)
N1—H1B \cdots O22 ⁱⁱ	0.90 (7)	2.55 (8)	3.290 (11)	141 (9)
N1—H1B \cdots O24 ⁱⁱ	0.90 (7)	2.40 (7)	3.272 (10)	166 (9)
N9—H9A \cdots O22 ⁱ	0.94 (4)	2.38 (6)	3.050 (8)	128 (5)
N12—H12A \cdots O26	0.92 (3)	2.36 (5)	3.046 (8)	132 (4)
N12—H12B \cdots O27 ⁱ	0.91 (5)	2.49 (5)	3.096 (8)	125 (4)
N20—H20B \cdots O26 ⁱⁱⁱ	0.91 (7)	2.56 (8)	3.246 (12)	133 (8)
N20—H20B \cdots O28 ⁱⁱⁱ	0.91 (7)	2.32 (8)	3.204 (11)	165 (8)
C8—H8B \cdots O24 ⁱⁱ	0.97	2.46	3.327 (9)	149
C10—H10A \cdots O24 ⁱⁱ	0.97	2.41	3.258 (8)	145
C10—H10B \cdots O22 ^{iv}	0.97	2.58	3.291 (9)	130
C11—H11A \cdots O27 ⁱ	0.97	2.56	3.147 (9)	119
C11—H11A \cdots O26 ^v	0.97	2.57	3.285 (9)	131
C11—H11B \cdots O28 ⁱⁱⁱ	0.97	2.41	3.249 (9)	145
C13—H13A \cdots O28 ⁱⁱⁱ	0.97	2.46	3.315 (10)	147

Symmetry codes: (i) $x, y + 1, z$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z$; (iii) $x - \frac{1}{2}, -y + \frac{1}{2}, z$; (iv) $x + \frac{1}{2}, -y + \frac{1}{2}, z$; (v) $x - \frac{1}{2}, -y + \frac{3}{2}, z$.

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS*; data reduction: *XSCANS*; program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2013*.

The authors thank the FCQ-UANL (project No. 03-6375-QMT-08-005) and PAICYT (Project No. IT164-09) for financial support. LAGR acknowledges a grant from the

CONACyT program "Dirección de Tesis entre la UANL y la University of Texas at Austin y/o Instituciones de Educación Superior de la ANUIES" (grant N.L.-2006-C09 32658).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: RK2415).

References

- Etter, M. C. (1990). *Acc. Chem. Res.* **23**, 120–126.
- Garza Rodríguez, L. Á. (2010). PhD thesis, Universidad Autónoma de Nuevo León, Mexico.
- Garza Rodríguez, L. Á., Bernès, S., Elizondo Martínez, P., Nájera Martínez, B. & Rodríguez de Luna, S. L. (2011). *Acta Cryst. E* **67**, o3235–o3236.
- Liu, Y.-F., Xia, H.-T., Wang, D.-Q., Yang, S.-P. & Meng, Y.-L. (2007). *Acta Cryst. E* **63**, o3836.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
- Rodríguez de Barbarán, C., Bernès, S., Nájera, B., Elizondo, P. & Cerdá, P. (2007). *Acta Cryst. E* **63**, o549–o550.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Siemens (1996). *XSCANS*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Yang, S.-P., Han, L.-J., Wang, D.-Q. & Xia, H.-T. (2007). *Acta Cryst. E* **63**, o3880.

supplementary materials

Acta Cryst. (2013). E69, o1643–o1644 [doi:10.1107/S1600536813027475]

N,N'-Bis(2-aminobenzyl)ethane-1,2-diaminium dinitrate

Luis Ángel Garza Rodríguez, Perla Elizondo Martínez, Sylvain Bernès, Blanca Nájera Martínez and Nancy Pérez Rodríguez

1. Comment

The title salt crystallized unexpectedly, when attempting the crystallization of a macrocyclic molecule, resulting from the Schiff condensation between 2,6-diacetylpyridine and *N,N'*-bis(2-aminobenzyl)ethane-1,2-diamine (Garza Rodríguez, 2010). The synthesis was carried out *via* a template reaction, using Mn²⁺ as metal center, and analytical data showed that the Mn²⁺ complex was formed, with nitrate as counter ions. However, this compound is almost insoluble in organic solvents, like MeOH, EtOH, acetone and ethyl acetate, impeding the preparation of single crystals. Only slight solubility was obtained in hot acetonitrile. Slow evaporation of MeCN over 3 weeks afforded a mixture of amorphous brown solids and colourless needle-shaped crystals. We assume that the brown solids should be a mixture of manganese oxides, resulting from the hydrolysis of the complex induced by trace amounts of water and dissolved O₂. Minutes amounts of H₃O⁺ are then released, which promote the formation, and finally the crystallization of the nitrate salt of the protonated amine.

Title compound (Fig. 1) crystallizes in a non-centrosymmetric space group, with all atoms placed in general positions. However, the dication (C₁₆H₂₄N₄)²⁺ presents a non-crystallographic inversion center, with the central aliphatic chain extended in the all-*trans* conformation. This conformation was previously obtained for the same cation crystallized as *p*-toluenesulfonate salt, although in that case, the dication was placed on a crystallographic inversion center (Garza Rodríguez *et al.*, 2011). The free amine, for which the X-ray structure is also known (Rodríguez de Barbarín *et al.*, 2007) has a different solid state conformation, although preserving the centrosymmetric character. For the nitrate salt reported here, departure from centrosymmetry is small, as reflected, for example, by the dihedral angle between benzene rings, limited to 3.3 (6)°.

Nitrate ions positions are determined by the formation of hydrogen bonds with ammonium NH₂⁺ groups in the cation. All N—H···O angles for these contacts are close to 180°, and H···O separations are in the range 1.87 (3) Å to 1.96 (3) Å. Each independent anion, N21 and N25, bridges two cations related by cell translation in the [0 1 0] direction, forming a R₄⁴(18) ring motif (Etter, 1990; Fig. 2). This arrangement seems actually to be common in crystal structures involving ammonium and nitrate species, and R₄⁴(18) motifs are also formed in salts closely related to the title compound, for example with *N,N'*-dibenzylethane-1,2-diammonium (Liu *et al.*, 2007) or *N,N'*-bis(4-chlorobenzyl)ethane-1,2-di-ammonium (Yang *et al.*, 2007). For such salts, the crystal structure is invariably based on edge-fused R₄⁴(18) rings, which afford a one-dimensional linear supramolecular structure. In the case of the title compound, chains run along the short *b* axis, and no significant interchain interactions are observed (Fig. 2).

2. Experimental

An amount of 2,6-diacetylpyridine (735 mg, 4.50 mmol) in ethanol (180 ml) was mixed with the templating reagent $Mn(NO_3)_2 \cdot xH_2O$ (1.130 g) and refluxed for 30 min. Then, N,N' -bis(2-aminobenzyl)ethane-1,2-diamine (1.302 g, 4.80 mmol, dissolved in 25 ml of ethanol) was slowly added, and the mixture further refluxed for 1 h. The resulting solid was filtered from hot ethanol, washed with cold ethanol, and dried under reduced pressure. A solution of the solid in hot CH_3CN was left to crystallize for 3 weeks. After this time, only one product was obtained as single crystals, in low yield, which was identified by X-ray diffraction as the title nitrate salt.

3. Refinement

H atoms for aromatic CH and methylene CH_2 groups were placed in idealized positions, and refined with C—H bond lengths fixed to 0.93 Å and 0.97 Å, respectively, and $U_{iso}(H) = 1.2U_{eq}(C)$. Amine and ammonium H atoms were found in a difference map, and refined freely, although the geometry for NH_2 group was restrained to sensible target values: bond lengths N—H were restrained to 0.90 (2) Å and $H \cdots H$ separations were restrained to 1.54 (3) Å. Isotropic displacement parameters were computed in this case as $U_{iso}(H) = 1.5U_{eq}(N)$.

Computing details

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS* (Siemens, 1996); data reduction: *XSCANS* (Siemens, 1996); program(s) used to solve structure: *SHELXS2013* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2013* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL2013* (Sheldrick, 2008).





Figure 1


Asymmetric unit of the title compound with the atom numbering scheme. Displacement ellipsoids for non-H atoms are presented at the 50% probability level. H atoms are shown as stick.

**Figure 2**

Part of the crystal structure of the title compound, showing N—H···O(nitrate) H bonds as dashed lines. Two supramolecular $R(18)$ -based chains are shown, which are related by the n glide plane perpendicular to [1 0 0]. No significant contacts are observed between chains.

N,N'-Bis(2-aminobenzyl)ethane-1,2-diaminium dinitrate

Crystal data



$M_r = 396.41$

Orthorhombic, $Pna2_1$

$a = 11.041 (5) \text{ \AA}$

$b = 5.760 (4) \text{ \AA}$

$c = 30.069 (13) \text{ \AA}$

$V = 1912.1 (18) \text{ \AA}^3$

$Z = 4$

$F(000) = 840$

$D_x = 1.377 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 75 reflections

$\theta = 4.9\text{--}11.7^\circ$

$\mu = 0.11 \text{ mm}^{-1}$

$T = 298\text{ K}$
Needle, colourless

$0.60 \times 0.20 \times 0.20\text{ mm}$

Data collection

Siemens P4
diffractometer
Radiation source: fine-focus sealed tube, FN4
Graphite monochromator
 $2\theta/\omega$ -scans
4371 measured reflections
2473 independent reflections
1501 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.053$
 $\theta_{\text{max}} = 25.5^\circ, \theta_{\text{min}} = 2.7^\circ$
 $h = -13 \rightarrow 13$
 $k = -3 \rightarrow 6$
 $l = -36 \rightarrow 36$
3 standard reflections every 97 reflections
intensity decay: 1.5%

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.061$
 $wR(F^2) = 0.180$
 $S = 1.61$
2473 reflections
278 parameters
13 restraints
0 constraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: difference Fourier map
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.06P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} < 0.001$
 $\Delta\rho_{\text{max}} = 0.34\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$
Extinction correction: *SHELXL2013* (Sheldrick, 2008), $F_c^* = kFc[1 + 0.001xFc^2\lambda^3/\sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.010 (2)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.9978 (7)	1.1615 (13)	0.1913 (3)	0.083 (2)
H1A	1.023 (9)	1.261 (14)	0.171 (2)	0.124*
H1B	1.038 (8)	1.148 (19)	0.2170 (18)	0.124*
C2	0.9002 (7)	1.0332 (13)	0.1770 (3)	0.0616 (19)
C3	0.8321 (9)	1.1154 (15)	0.1408 (3)	0.077 (2)
H3A	0.8541	1.2543	0.1272	0.092*
C4	0.7348 (9)	0.9958 (16)	0.1251 (3)	0.087 (3)
H4A	0.6902	1.0542	0.1014	0.104*
C5	0.7024 (8)	0.7886 (18)	0.1444 (3)	0.080 (2)
H5A	0.6364	0.7049	0.1339	0.096*
C6	0.7699 (7)	0.7072 (16)	0.1797 (2)	0.067 (2)
H6A	0.7485	0.5667	0.1928	0.080*
C7	0.8672 (7)	0.8267 (13)	0.1961 (2)	0.0564 (18)
C8	0.9347 (6)	0.7329 (14)	0.2354 (2)	0.0545 (18)
H8A	0.9438	0.5662	0.2322	0.065*
H8B	1.0150	0.8010	0.2363	0.065*
N9	0.8705 (4)	0.7843 (10)	0.27754 (19)	0.0485 (13)
H9A	0.846 (6)	0.937 (5)	0.284 (3)	0.073*
H9B	0.795 (4)	0.706 (10)	0.276 (3)	0.073*
C10	0.9427 (5)	0.7325 (13)	0.3181 (2)	0.0492 (15)
H10A	1.0124	0.8346	0.3195	0.059*
H10B	0.9715	0.5734	0.3171	0.059*

C11	0.8645 (5)	0.7674 (13)	0.3583 (2)	0.0474 (14)
H11A	0.8372	0.9274	0.3597	0.057*
H11B	0.7937	0.6679	0.3566	0.057*
N12	0.9349 (4)	0.7109 (11)	0.39831 (19)	0.0489 (13)
H12A	0.960 (5)	0.559 (5)	0.398 (3)	0.073*
H12B	0.999 (4)	0.810 (9)	0.397 (3)	0.073*
C13	0.8723 (7)	0.7589 (13)	0.4415 (2)	0.0559 (18)
H13A	0.7905	0.6976	0.4406	0.067*
H13B	0.8674	0.9252	0.4463	0.067*
C14	0.9406 (7)	0.6483 (14)	0.4791 (2)	0.056 (2)
C15	1.0413 (8)	0.7526 (16)	0.4955 (2)	0.073 (2)
H15A	1.0652	0.8941	0.4835	0.087*
C16	1.1096 (9)	0.658 (2)	0.5293 (3)	0.093 (3)
H16A	1.1798	0.7300	0.5392	0.112*
C17	1.0704 (10)	0.451 (2)	0.5479 (3)	0.094 (3)
H17A	1.1137	0.3844	0.5711	0.113*
C18	0.9691 (9)	0.3471 (17)	0.5323 (3)	0.085 (3)
H18A	0.9439	0.2083	0.5451	0.102*
C19	0.9016 (7)	0.4411 (14)	0.4977 (3)	0.066 (2)
N20	0.7978 (8)	0.3323 (14)	0.4831 (3)	0.086 (2)
H20A	0.786 (10)	0.185 (8)	0.493 (3)	0.128*
H20B	0.764 (9)	0.363 (17)	0.4563 (19)	0.128*
N21	0.7142 (5)	0.2743 (11)	0.2797 (2)	0.0550 (14)
O22	0.6509 (5)	0.1048 (10)	0.2738 (2)	0.093 (2)
O23	0.8219 (4)	0.2579 (8)	0.28670 (19)	0.0729 (15)
O24	0.6674 (4)	0.4708 (9)	0.27685 (19)	0.0735 (15)
N25	1.0910 (5)	0.2211 (10)	0.3951 (2)	0.0556 (14)
O26	1.1542 (5)	0.3900 (10)	0.3991 (3)	0.099 (2)
O27	0.9825 (4)	0.2383 (9)	0.38897 (19)	0.0702 (16)
O28	1.1372 (5)	0.0215 (9)	0.3989 (2)	0.0750 (14)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.097 (5)	0.055 (5)	0.095 (5)	-0.021 (4)	-0.007 (4)	0.007 (4)
C2	0.077 (5)	0.041 (4)	0.066 (4)	-0.003 (4)	0.006 (4)	0.001 (4)
C3	0.110 (7)	0.055 (5)	0.066 (4)	-0.006 (5)	-0.004 (5)	0.010 (4)
C4	0.104 (7)	0.085 (7)	0.071 (5)	-0.002 (7)	-0.015 (5)	0.011 (5)
C5	0.074 (5)	0.100 (7)	0.067 (4)	-0.016 (5)	-0.012 (4)	-0.002 (5)
C6	0.063 (4)	0.069 (6)	0.068 (5)	-0.007 (4)	-0.006 (4)	0.000 (4)
C7	0.061 (4)	0.051 (5)	0.057 (4)	0.001 (4)	0.005 (3)	0.006 (4)
C8	0.060 (4)	0.047 (4)	0.057 (4)	0.003 (4)	0.009 (3)	0.002 (4)
N9	0.051 (3)	0.036 (3)	0.058 (3)	0.005 (3)	-0.006 (3)	-0.002 (3)
C10	0.048 (4)	0.039 (4)	0.061 (4)	0.000 (3)	-0.003 (3)	0.002 (4)
C11	0.048 (3)	0.040 (4)	0.054 (3)	0.003 (3)	-0.005 (3)	0.003 (4)
N12	0.053 (3)	0.037 (3)	0.056 (3)	0.006 (3)	0.003 (3)	0.003 (3)
C13	0.064 (4)	0.042 (4)	0.062 (4)	0.001 (4)	0.005 (3)	0.008 (4)
C14	0.066 (5)	0.047 (5)	0.054 (4)	0.000 (4)	0.001 (3)	-0.004 (4)
C15	0.080 (5)	0.079 (6)	0.059 (4)	-0.002 (5)	-0.005 (4)	-0.007 (5)
C16	0.089 (6)	0.119 (9)	0.071 (5)	-0.003 (6)	-0.009 (5)	-0.016 (6)

C17	0.102 (7)	0.120 (9)	0.060 (5)	0.040 (7)	-0.005 (5)	-0.006 (6)
C18	0.114 (7)	0.077 (7)	0.062 (5)	0.026 (6)	0.015 (5)	0.014 (5)
C19	0.082 (5)	0.050 (5)	0.065 (4)	0.010 (4)	0.012 (4)	-0.002 (4)
N20	0.096 (6)	0.057 (5)	0.103 (5)	-0.015 (5)	0.002 (5)	0.009 (4)
N21	0.050 (3)	0.041 (4)	0.074 (3)	0.005 (3)	0.001 (3)	0.005 (4)
O22	0.077 (4)	0.049 (3)	0.151 (6)	-0.016 (3)	-0.005 (4)	-0.003 (4)
O23	0.051 (3)	0.043 (3)	0.125 (4)	0.000 (3)	-0.014 (3)	-0.004 (4)
O24	0.063 (3)	0.047 (3)	0.111 (4)	0.015 (3)	0.002 (3)	0.004 (3)
N25	0.054 (3)	0.037 (4)	0.075 (3)	0.005 (3)	0.003 (3)	0.001 (4)
O26	0.074 (4)	0.050 (4)	0.174 (6)	-0.023 (3)	-0.001 (4)	0.005 (5)
O27	0.055 (3)	0.039 (3)	0.116 (4)	0.004 (3)	-0.013 (3)	0.008 (4)
O28	0.067 (3)	0.045 (3)	0.113 (4)	0.014 (3)	0.001 (3)	0.005 (3)

Geometric parameters (\AA , $\text{^{\circ}}$)

N1—C2	1.376 (10)	N12—C13	1.498 (9)
N1—H1A	0.89 (2)	N12—H12A	0.92 (2)
N1—H1B	0.90 (2)	N12—H12B	0.91 (2)
C2—C7	1.371 (9)	C13—C14	1.500 (10)
C2—C3	1.405 (11)	C13—H13A	0.9700
C3—C4	1.360 (12)	C13—H13B	0.9700
C3—H3A	0.9300	C14—C15	1.356 (11)
C4—C5	1.374 (12)	C14—C19	1.386 (11)
C4—H4A	0.9300	C15—C16	1.378 (12)
C5—C6	1.379 (11)	C15—H15A	0.9300
C5—H5A	0.9300	C16—C17	1.382 (15)
C6—C7	1.368 (10)	C16—H16A	0.9300
C6—H6A	0.9300	C17—C18	1.354 (12)
C7—C8	1.497 (10)	C17—H17A	0.9300
C8—N9	1.482 (8)	C18—C19	1.389 (12)
C8—H8A	0.9700	C18—H18A	0.9300
C8—H8B	0.9700	C19—N20	1.378 (11)
N9—C10	1.486 (8)	N20—H20A	0.91 (2)
N9—H9A	0.94 (2)	N20—H20B	0.91 (2)
N9—H9B	0.95 (2)	N21—O23	1.212 (6)
C10—C11	1.501 (8)	N21—O22	1.213 (7)
C10—H10A	0.9700	N21—O24	1.247 (7)
C10—H10B	0.9700	N25—O26	1.203 (7)
C11—N12	1.468 (9)	N25—O27	1.215 (7)
C11—H11A	0.9700	N25—O28	1.263 (7)
C11—H11B	0.9700		
C2—N1—H1A	112 (6)	N12—C11—H11B	109.9
C2—N1—H1B	128 (6)	C10—C11—H11B	109.9
H1A—N1—H1B	120 (5)	H11A—C11—H11B	108.3
C7—C2—N1	122.8 (7)	C11—N12—C13	115.1 (5)
C7—C2—C3	118.4 (7)	C11—N12—H12A	112 (5)
N1—C2—C3	118.8 (8)	C13—N12—H12A	109 (5)
C4—C3—C2	121.3 (8)	C11—N12—H12B	104 (5)
C4—C3—H3A	119.3	C13—N12—H12B	106 (5)

C2—C3—H3A	119.3	H12A—N12—H12B	111 (4)
C3—C4—C5	120.0 (8)	N12—C13—C14	110.1 (6)
C3—C4—H4A	120.0	N12—C13—H13A	109.6
C5—C4—H4A	120.0	C14—C13—H13A	109.6
C4—C5—C6	118.7 (9)	N12—C13—H13B	109.6
C4—C5—H5A	120.7	C14—C13—H13B	109.6
C6—C5—H5A	120.7	H13A—C13—H13B	108.2
C7—C6—C5	122.0 (9)	C15—C14—C19	119.3 (8)
C7—C6—H6A	119.0	C15—C14—C13	119.9 (7)
C5—C6—H6A	119.0	C19—C14—C13	120.8 (7)
C6—C7—C2	119.6 (7)	C14—C15—C16	122.7 (9)
C6—C7—C8	119.6 (7)	C14—C15—H15A	118.6
C2—C7—C8	120.8 (7)	C16—C15—H15A	118.6
N9—C8—C7	111.4 (5)	C15—C16—C17	117.9 (10)
N9—C8—H8A	109.3	C15—C16—H16A	121.1
C7—C8—H8A	109.3	C17—C16—H16A	121.1
N9—C8—H8B	109.3	C18—C17—C16	120.0 (9)
C7—C8—H8B	109.3	C18—C17—H17A	120.0
H8A—C8—H8B	108.0	C16—C17—H17A	120.0
C8—N9—C10	113.9 (5)	C17—C18—C19	121.9 (9)
C8—N9—H9A	120 (5)	C17—C18—H18A	119.0
C10—N9—H9A	100 (5)	C19—C18—H18A	119.0
C8—N9—H9B	107 (5)	N20—C19—C14	121.5 (8)
C10—N9—H9B	114 (5)	N20—C19—C18	120.4 (9)
H9A—N9—H9B	102 (4)	C14—C19—C18	118.1 (8)
N9—C10—C11	109.0 (4)	C19—N20—H20A	116 (6)
N9—C10—H10A	109.9	C19—N20—H20B	122 (6)
C11—C10—H10A	109.9	H20A—N20—H20B	115 (5)
N9—C10—H10B	109.9	O23—N21—O22	121.8 (6)
C11—C10—H10B	109.9	O23—N21—O24	119.3 (6)
H10A—C10—H10B	108.3	O22—N21—O24	118.8 (6)
N12—C11—C10	109.0 (4)	O26—N25—O27	121.4 (6)
N12—C11—H11A	109.9	O26—N25—O28	119.5 (6)
C10—C11—H11A	109.9	O27—N25—O28	119.1 (6)
C7—C2—C3—C4	0.8 (12)	C10—C11—N12—C13	174.5 (5)
N1—C2—C3—C4	179.9 (9)	C11—N12—C13—C14	167.8 (7)
C2—C3—C4—C5	-1.0 (14)	N12—C13—C14—C15	79.3 (9)
C3—C4—C5—C6	0.4 (14)	N12—C13—C14—C19	-101.3 (7)
C4—C5—C6—C7	0.2 (13)	C19—C14—C15—C16	2.1 (12)
C5—C6—C7—C2	-0.4 (12)	C13—C14—C15—C16	-178.5 (7)
C5—C6—C7—C8	178.2 (7)	C14—C15—C16—C17	-2.2 (13)
N1—C2—C7—C6	-179.2 (7)	C15—C16—C17—C18	1.2 (13)
C3—C2—C7—C6	-0.1 (11)	C16—C17—C18—C19	0.0 (13)
N1—C2—C7—C8	2.2 (11)	C15—C14—C19—N20	177.9 (8)
C3—C2—C7—C8	-178.7 (7)	C13—C14—C19—N20	-1.6 (11)
C6—C7—C8—N9	-79.6 (9)	C15—C14—C19—C18	-0.8 (11)
C2—C7—C8—N9	99.1 (7)	C13—C14—C19—C18	179.8 (7)
C7—C8—N9—C10	-169.9 (6)	C17—C18—C19—N20	-178.9 (8)

C8—N9—C10—C11	−174.2 (6)	C17—C18—C19—C14	−0.2 (12)
N9—C10—C11—N12	178.7 (7)		

Hydrogen-bond geometry (Å, °)

D—H···A	D—H	H···A	D···A	D—H···A
N9—H9A···O23 ⁱ	0.94 (4)	1.87 (3)	2.794 (8)	167 (7)
N9—H9B···O24	0.95 (5)	1.96 (5)	2.879 (7)	164 (5)
N12—H12A···O27	0.92 (3)	1.88 (3)	2.787 (8)	168 (8)
N12—H12B···O28 ⁱ	0.91 (5)	1.95 (5)	2.862 (8)	176 (9)
N1—H1B···O22 ⁱⁱ	0.90 (7)	2.55 (8)	3.290 (11)	141 (9)
N1—H1B···O24 ⁱⁱ	0.90 (7)	2.40 (7)	3.272 (10)	166 (9)
N9—H9A···O22 ⁱ	0.94 (4)	2.38 (6)	3.050 (8)	128 (5)
N12—H12A···O26	0.92 (3)	2.36 (5)	3.046 (8)	132 (4)
N12—H12B···O27 ⁱ	0.91 (5)	2.49 (5)	3.096 (8)	125 (4)
N20—H20B···O26 ⁱⁱⁱ	0.91 (7)	2.56 (8)	3.246 (12)	133 (8)
N20—H20B···O28 ⁱⁱⁱ	0.91 (7)	2.32 (8)	3.204 (11)	165 (8)
C8—H8B···O24 ⁱⁱ	0.97	2.46	3.327 (9)	149
C10—H10A···O24 ⁱⁱ	0.97	2.41	3.258 (8)	145
C10—H10B···O22 ^{iv}	0.97	2.58	3.291 (9)	130
C11—H11A···O27 ⁱ	0.97	2.56	3.147 (9)	119
C11—H11A···O26 ^v	0.97	2.57	3.285 (9)	131
C11—H11B···O28 ⁱⁱⁱ	0.97	2.41	3.249 (9)	145
C13—H13A···O28 ⁱⁱⁱ	0.97	2.46	3.315 (10)	147

Symmetry codes: (i) $x, y+1, z$; (ii) $x+1/2, -y+3/2, z$; (iii) $x-1/2, -y+1/2, z$; (iv) $x+1/2, -y+1/2, z$; (v) $x-1/2, -y+3/2, z$.