

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/221761135>

4-[(E)-(4-Fluorobenzylidene)amino]benzoic acid

Article in *Acta Crystallographica Section E Structure Reports Online* · January 2012

DOI: 10.1107/S1600536811052275 · Source: PubMed

CITATION

1

READS

45

5 authors, including:



[Victor Manuel Jimenez Perez](#)
Autonomous University of Nuevo León

7 PUBLICATIONS 35 CITATIONS

[SEE PROFILE](#)



[Noemí Waksman de Torres](#)
Autonomous University of Nuevo León

96 PUBLICATIONS 781 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Study in vivo and in vitro antiviral activity of *Jatropha dioica*. [View project](#)



Leucophyllum frutescens [View project](#)

4-[(E)-(4-Fluorobenzylidene)amino]-benzoic acid

Blanca M. Muñoz-Flores,^{a*} Víctor M. Jiménez Pérez,^a Rosa L. Santillan,^b María Eugenia Ochoa^b and Noemí Waksman^c

^aFacultad de Ciencias Químicas, Universidad Autónoma de Nuevo León, San Nicolás de los Garza, 66451 Nuevo León, Mexico, ^bDepartamento de Química, Centro de Investigación y de Estudios Avanzados, del Instituto Politécnico Nacional, Apartado Postal 14-740, 07000 México, DF, Mexico, and ^cDepartamento de Química Analítica Facultad de Medicina, Universidad Autónoma de Nuevo León, León PO Box 2316, 64841 Nuevo León, Mexico
Correspondence e-mail: blanca.munozfl@uanl.edu.mx


Received 16 November 2011; accepted 4 December 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$; R factor = 0.062; wR factor = 0.191; data-to-parameter ratio = 12.1.

In the title compound, $\text{C}_{14}\text{H}_{10}\text{FNO}_2$, the benzene rings make a dihedral angle of $57.50(13)^\circ$, and the molecule has an *E* configuration about the $\text{C}=\text{N}$ bond. In the crystal, molecules are linked *via* pairs of $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds, forming inversion dimers.

Related literature

For the synthesis, properties and uses of 4-(benzylidene)amino)benzoic acid, see: Borisova *et al.* (2007); Schiff (1864); Innocenzi & Lebeau (2005); Muñoz-Flores *et al.* (2008).



Experimental

Crystal data

$\text{C}_{14}\text{H}_{10}\text{FNO}_2$	$c = 17.2874(8)\text{ \AA}$
$M_r = 243.24$	$\beta = 105.833(2)^\circ$
Monoclinic, $P2_1/n$	$V = 1148.99(8)\text{ \AA}^3$
$a = 12.2787(5)\text{ \AA}$	$Z = 4$
$b = 5.6264(2)\text{ \AA}$	Mo $K\alpha$ radiation

$\mu = 0.11\text{ mm}^{-1}$
 $T = 293\text{ K}$

$0.23 \times 0.2 \times 0.15\text{ mm}$

Data collection

Nonius KappaCCD diffractometer
9975 measured reflections
2002 independent reflections
1226 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.186$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.062$
 $wR(F^2) = 0.191$
 $S = 1.04$
2002 reflections
166 parameters
1 restraint
H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.18\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.21\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H ₂ —O1 ⁱ	0.85 (1)	1.79 (2)	2.601 (3)	159 (4)

Symmetry code: (i) $-x + 1, -y + 1, -z + 1$.

Data collection: *COLLECT* (Nonius, 1998); cell refinement: *XSCANS* (Bruker, 2000); data reduction: *XSCANS*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2006); software used to prepare material for publication: *SHELXL97*.

The authors thank the Programa de Mejoramiento del Profesorado (PROMEP; grant No. 103.5/11/4330) and the Consejo Nacional de Ciencia y Tecnología (CONACYT; grant No. 82605) for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: FJ2486).

References

- Borisova, N. E., Reshetova, M. D. & Ustynguk, Y. A. (2007). *Chem. Rev.* **107**, 46–79.
- Bruker (2000). *XSCANS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Innocenzi, P. & Lebeau, B. (2005). *J. Mater. Chem.* **15**, 3821–3831.
- Macrae, C. F., Edgington, P. R., McCabe, P., Pidcock, E., Shields, G. P., Taylor, R., Towler, M. & van de Streek, J. (2006). *J. Appl. Cryst.* **39**, 453–457.
- Muñoz-Flores, B. M., Santillan, R., Rodríguez, M., Méndez, J. M., Romero, M., Farfán, N., Lacroix, P. G., Nakatani, K., Ramos-Ortíz, G. & Maldonado, J. L. (2008). *J. Organomet. Chem.* **693**, 1321–1334.
- Nonius (1998). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Schiff, H. (1864). *Annalen*, **131**, 118–119.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o175 [doi:10.1107/S1600536811052275]

4-[*(E*)-(4-Fluorobenzylidene)amino]benzoic acid

B. M. Muñoz-Flores, V. M. J. Pérez, R. L. Santillan, M. E. Ochoa and N. Waksman

Comment

Hugo Josef Schiff discovered the condensation a primary amine and a carbonyl compounds (Schiff 1864) afford the corresponding azomethine group. This organic compounds show an important synthetic advantages such as: high yields, simple synthetic route, short-time reactions, and easy isolation. Schiff base compounds have been of great importance in coordination chemistry (Borisova *et al.* 2007) due to the lone pair of electrons in an sp^2 hybridized orbital of nitrogen atom of the azomethine group. On the other hand, it has been reported that organic compounds containing donor and acceptor groups linked through to pi-system delocalized exhibit promising nonlinear optical properties (Innocenzi & Lebeau 2005). We have been focused our attention on synthesis of *push-pull* organic molecules with no linear optical potential properties (Muñoz-Flores *et al.* 2008). In continuous with our research, we synthesized the title compound (*E*)-4-(4-fluorobenzylideneamino)benzoic acid by condensation of 4-fluorobenzaldehyde and 4-aminobenzoic acid. In the present article, the crystal structure of (I) is being reported as shown in Fig 1. The compound, (*E*)-4-(4-fluorobenzylideneamino)benzoic acid ($C_{14}H_{10}N_2O_2F$), displays C_1 symmetry. The aromatic rings are not in the same plane, with a dihedral angle of 15.59° between mean planes. The carboxylic group represents a delocalized system with C(14)—O(1) and C(14)—O(2) bond lengths are 1.275 (4) and 1.293 (4) Å, respectively. The azomethine group are in the same plane as the monofluorated ring, probably because the short contact between the C(5)—H(5)…N(1) 2.633 (4) Å, [$\langle C—H\cdots N \rangle$: 96.8°]. The intermolecular O(2)—H(2)…O(1) [2.621 (5) Å (angle O-H…O: 156.30°], hydrogen bond form a dimer with a inversion center as shown in Fig 2.

Experimental

A solution of 4-fluorobenzaldehyde (0.5 g, 4 mmol) and 4-aminobenzoic acid (0.55 g, 4 mmol) in benzene (50 ml) was heated under reflux for 6 h, with a Dean-Stark apparatus used for the azeotropic removal of water and allowed to cool to room temperature. Removal of solvent yielded a pale yellow solid, which was recrystallized from hot benzene (10 ml). Yield: 0.74 g 76%. *M. p.* 191 °C. *1H NMR* (400.13 MHz, MeOD): δ = 4.9 (bs, 1H, OH), 6.62 (d, 3J = 8.4 Hz, 2H, H-11/H-12), 6.80 (d, 3J = 8.4 Hz, 2H, H-9/H-13), 7.10 (t, 3J = 8.4 Hz, 2H, H-2/H-4), 8.05 (d, 3J = 78.4 Hz, 2H, H-1/H-5), 8.53 (s, 1H, H-7). *MS* (DIP 20 eV) for $C_{14}H_{10}N_2O_2F$ (f. w: 243.24 g/mol) *m/z* (%): 243 (100) [M^+], 226 (3) [$M^+ - H_2O$], 198 (3) [$M^+ - CO_2$], 137 (4) [$M^+ - FC_6H_8$], 121 [$M^+ - C_7H_6O_2$].

Refinement

All C-bonded H atoms were placed in calculated positions and refined as riding to their carrier atoms, with bond lengths fixed to 0.93 (aromatic CH). Isotropic displacement parameters for H atoms were calculated as $U_{iso}(H) = 1.5U_{eq}$ (carrier atom).

supplementary materials

Figures




Fig. 1. The molecular structure of (I), with atom labels and 50% probability displacement ellipsoids for non-H atoms.




Fig. 2. Intermolecular interaction *via* hydrogen bonds.

4-[*(E*)-(4-Fluorobenzylidene)amino]benzoic acid

Crystal data

C ₁₄ H ₁₀ FNO ₂	<i>F</i> (000) = 504
<i>M_r</i> = 243.24	<i>D_x</i> = 1.406 Mg m ⁻³
Monoclinic, <i>P2₁/n</i>	Melting point: 464 K
Hall symbol: -P 2yn	Mo <i>Kα</i> radiation, λ = 0.71073 Å
<i>a</i> = 12.2787 (5) Å	Cell parameters from 4962 reflections
<i>b</i> = 5.6264 (2) Å	θ = 2.9–27.5°
<i>c</i> = 17.2874 (8) Å	μ = 0.11 mm ⁻¹
β = 105.833 (2)°	<i>T</i> = 293 K
<i>V</i> = 1148.99 (8) Å ³	Prism, yellow
<i>Z</i> = 4	0.23 × 0.2 × 0.15 mm

Data collection

Nonius KappaCCD diffractometer	1226 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube graphite	R_{int} = 0.186
CCD rotation images, thick slices scans	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 3.7^\circ$
9975 measured reflections	$h = -14 \rightarrow 14$
2002 independent reflections	$k = -6 \rightarrow 6$
	$l = -17 \rightarrow 20$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)]$ = 0.062	Hydrogen site location: inferred from neighbouring sites
$wR(F^2)$ = 0.191	H atoms treated by a mixture of independent and constrained refinement
S = 1.04	$w = 1/[\sigma^2(F_o^2) + (0.0979P)^2 + 0.0699P]$ where $P = (F_o^2 + 2F_c^2)/3$
2002 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
166 parameters	$\Delta\rho_{\text{max}} = 0.18 \text{ e \AA}^{-3}$

1 restraint

$$\Delta\rho_{\min} = -0.21 \text{ e \AA}^{-3}$$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7423 (2)	0.3339 (5)	-0.03261 (18)	0.0515 (8)
H1	0.7623	0.4696	-0.0012	0.062*
C2	0.7740 (2)	0.3115 (5)	-0.10261 (19)	0.0565 (8)
H2	0.8169	0.4284	-0.1185	0.068*
C3	0.7407 (3)	0.1119 (5)	-0.14822 (19)	0.0557 (8)
C4	0.6800 (2)	-0.0683 (5)	-0.12777 (19)	0.0567 (8)
H4	0.6588	-0.2005	-0.1607	0.068*
C5	0.6511 (2)	-0.0474 (5)	-0.05617 (18)	0.0528 (7)
H5	0.6116	-0.1698	-0.0397	0.063*
C6	0.6803 (2)	0.1547 (4)	-0.00823 (17)	0.0479 (7)
C7	0.6428 (2)	0.1789 (5)	0.06460 (17)	0.0506 (7)
H7	0.6186	0.0437	0.0861	0.061*
C8	0.6104 (2)	0.3845 (4)	0.17270 (17)	0.0452 (7)
C9	0.5490 (2)	0.5794 (4)	0.1872 (2)	0.0563 (8)
H9	0.5247	0.6939	0.1474	0.068*
C10	0.5237 (2)	0.6054 (5)	0.25934 (19)	0.0534 (8)
H10	0.4824	0.7367	0.2677	0.064*
C11	0.5597 (2)	0.4361 (4)	0.32045 (17)	0.0482 (7)
C12	0.6204 (2)	0.2400 (5)	0.30522 (19)	0.0554 (8)
H12	0.6446	0.1251	0.3449	0.066*
C13	0.6449 (2)	0.2136 (5)	0.23337 (18)	0.0546 (8)
H13	0.6849	0.0808	0.2246	0.065*
C14	0.5353 (2)	0.4651 (5)	0.39801 (18)	0.0523 (8)
F1	0.77081 (18)	0.0920 (3)	-0.21809 (12)	0.0827 (7)
N1	0.64180 (18)	0.3768 (4)	0.10000 (15)	0.0530 (7)
O1	0.56240 (18)	0.3041 (4)	0.45179 (13)	0.0679 (7)
O2	0.48502 (19)	0.6570 (4)	0.41100 (13)	0.0675 (7)
H	0.487 (3)	0.678 (7)	0.4597 (9)	0.101*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0568 (16)	0.0450 (14)	0.0523 (19)	-0.0016 (12)	0.0142 (13)	-0.0017 (12)
C2	0.0623 (17)	0.0512 (15)	0.061 (2)	0.0016 (13)	0.0245 (15)	0.0098 (14)
C3	0.0649 (18)	0.0582 (17)	0.0471 (19)	0.0168 (14)	0.0206 (14)	0.0037 (14)
C4	0.0639 (17)	0.0483 (15)	0.058 (2)	0.0042 (13)	0.0177 (15)	-0.0075 (14)
C5	0.0526 (15)	0.0457 (14)	0.062 (2)	-0.0007 (12)	0.0191 (13)	0.0006 (14)
C6	0.0455 (14)	0.0456 (14)	0.0520 (19)	0.0034 (11)	0.0122 (13)	0.0026 (12)
C7	0.0508 (15)	0.0520 (15)	0.0516 (18)	-0.0003 (12)	0.0185 (13)	0.0061 (13)
C8	0.0433 (14)	0.0493 (15)	0.0436 (17)	-0.0052 (11)	0.0131 (12)	-0.0007 (12)
C9	0.0569 (16)	0.0434 (14)	0.066 (2)	0.0042 (12)	0.0116 (14)	0.0095 (13)
C10	0.0558 (16)	0.0470 (14)	0.058 (2)	0.0056 (12)	0.0165 (14)	0.0001 (13)
C11	0.0478 (14)	0.0478 (14)	0.0477 (18)	0.0006 (11)	0.0106 (12)	0.0016 (13)

supplementary materials

C12	0.0628 (17)	0.0532 (16)	0.0474 (18)	0.0097 (13)	0.0106 (14)	0.0066 (13)
C13	0.0551 (16)	0.0503 (15)	0.056 (2)	0.0109 (12)	0.0112 (14)	0.0044 (13)
C14	0.0467 (14)	0.0511 (15)	0.056 (2)	0.0011 (13)	0.0085 (12)	-0.0023 (14)
F1	0.1162 (16)	0.0737 (12)	0.0682 (14)	0.0116 (11)	0.0420 (12)	0.0015 (10)
N1	0.0532 (14)	0.0494 (13)	0.0564 (16)	0.0016 (10)	0.0150 (11)	0.0027 (11)
O1	0.0832 (15)	0.0652 (13)	0.0558 (15)	0.0141 (11)	0.0198 (11)	0.0101 (10)
O2	0.0804 (15)	0.0660 (13)	0.0549 (15)	0.0158 (10)	0.0165 (12)	-0.0040 (11)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.375 (4)	C8—C13	1.400 (4)
C1—C6	1.396 (4)	C8—N1	1.413 (4)
C1—H1	0.9300	C9—C10	1.372 (4)
C2—C3	1.369 (4)	C9—H9	0.9300
C2—H2	0.9300	C10—C11	1.402 (4)
C3—C4	1.360 (4)	C10—H10	0.9300
C3—F1	1.361 (4)	C11—C12	1.396 (4)
C4—C5	1.383 (4)	C11—C14	1.460 (4)
C4—H4	0.9300	C12—C13	1.363 (4)
C5—C6	1.395 (4)	C12—H12	0.9300
C5—H5	0.9300	C13—H13	0.9300
C6—C7	1.460 (4)	C14—O1	1.275 (4)
C7—N1	1.272 (3)	C14—O2	1.294 (3)
C7—H7	0.9300	O2—H	0.845 (10)
C8—C9	1.392 (4)		
C2—C1—C6	120.6 (3)	C9—C8—N1	118.5 (2)
C2—C1—H1	119.7	C13—C8—N1	123.0 (2)
C6—C1—H1	119.7	C10—C9—C8	121.0 (3)
C3—C2—C1	118.0 (3)	C10—C9—H9	119.5
C3—C2—H2	121.0	C8—C9—H9	119.5
C1—C2—H2	121.0	C9—C10—C11	120.7 (3)
C4—C3—F1	118.0 (3)	C9—C10—H10	119.7
C4—C3—C2	124.1 (3)	C11—C10—H10	119.7
F1—C3—C2	117.9 (3)	C12—C11—C10	118.0 (3)
C3—C4—C5	117.5 (3)	C12—C11—C14	121.0 (2)
C3—C4—H4	121.2	C10—C11—C14	121.0 (2)
C5—C4—H4	121.2	C13—C12—C11	121.3 (3)
C4—C5—C6	120.9 (3)	C13—C12—H12	119.4
C4—C5—H5	119.5	C11—C12—H12	119.4
C6—C5—H5	119.5	C12—C13—C8	120.7 (3)
C5—C6—C1	118.8 (3)	C12—C13—H13	119.6
C5—C6—C7	119.9 (2)	C8—C13—H13	119.6
C1—C6—C7	121.3 (2)	O1—C14—O2	120.6 (3)
N1—C7—C6	122.9 (2)	O1—C14—C11	120.8 (2)
N1—C7—H7	118.5	O2—C14—C11	118.7 (3)
C6—C7—H7	118.5	C7—N1—C8	119.7 (2)
C9—C8—C13	118.3 (3)	C14—O2—H	114 (3)
C6—C1—C2—C3	1.6 (4)	C9—C10—C11—C12	-0.8 (4)
C1—C2—C3—C4	-1.4 (4)	C9—C10—C11—C14	178.6 (2)

C1—C2—C3—F1	179.0 (2)	C10—C11—C12—C13	0.4 (4)
F1—C3—C4—C5	179.2 (2)	C14—C11—C12—C13	-179.0 (3)
C2—C3—C4—C5	-0.3 (4)	C11—C12—C13—C8	0.5 (4)
C3—C4—C5—C6	1.9 (4)	C9—C8—C13—C12	-1.0 (4)
C4—C5—C6—C1	-1.8 (4)	N1—C8—C13—C12	174.3 (2)
C4—C5—C6—C7	176.2 (2)	C12—C11—C14—O1	-4.7 (4)
C2—C1—C6—C5	0.0 (4)	C10—C11—C14—O1	175.9 (2)
C2—C1—C6—C7	-178.0 (2)	C12—C11—C14—O2	175.7 (2)
C5—C6—C7—N1	-162.4 (3)	C10—C11—C14—O2	-3.7 (4)
C1—C6—C7—N1	15.6 (4)	C6—C7—N1—C8	-176.5 (2)
C13—C8—C9—C10	0.6 (4)	C9—C8—N1—C7	-143.9 (3)
N1—C8—C9—C10	-174.9 (2)	C13—C8—N1—C7	40.8 (4)
C8—C9—C10—C11	0.3 (4)		


Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O2—H···O1 ⁱ	0.85 (1)	1.79 (2)	2.601 (3)	159 (4)

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

supplementary materials

Fig. 1



supplementary materials

Fig. 2

