

- errors: modeling the origins of genomic instability. *Oncogene* 2002;21:6241-8.
46. Gisselsson D, Pettersson L, Hoglund M, Heidenblad M, Gorunova L, Wiegant J, et al. Chromosomal breakage fusion-bridge events cause genetic intratumor heterogeneity. *Proc Natl Acad Sci USA* 2002;97:5357-62.
 47. McClintock B. The stability of broken ends of chromosomes in *Zea mays*. *Genetics* 1940;26:234-82.
 48. Plug-Demaggio AW, McDougall JK. The human papillomavirus type 16 E6 oncogene induces premature mitotic chromosome segregation. *Oncogene* 2002;21:7507-13.
 49. Williams VM, Filippova M, Filippov V, Payne KP, Duerksen-Hughesa P. Human papillomavirus Type 16 E6* induces oxidative stress and DNA damage. *J Virol* 2014;88: 6751-61.
 50. Marullo R, Werner E, Zhang H, Chen GZ, Shin DM, Doetsch PW. HPV16 E6 and E7 proteins induce a chronic oxidative stress response via NOX2 that causes genomic instability and increased susceptibility to DNA damage in head and neck cancer cells. *Carcinogenesis* 2015;36:1397-406.
 51. Cortés-Gutiérrez EI, Dávila-Rodríguez MI, Vargas-Villarreal J, Hernández-Garza F, Cerda-Flores RM. Association between human papilloma virus-type infections with micronuclei frequencies. *Prague Med Rep* 2010;111:35-41.
 52. Maser RS, DePinho RA. Keeping telomerase in its place. *Nat Med* 2002;8:934-6.
 53. Artandi SE, Chang S, Lee SL, Alson S, Gottlieb GJ, Chin L, et al. Telomere dysfunction promotes non-reciprocal translocations and epithelial cancers in mice. *Nature* 2000;10:641-5.
 54. Rudolph KL, Millard M, Bosenberg MW, DePinho RA. Telomere dysfunction and evolution of intestinal carcinoma in mice and humans. *Nat Genet* 2001;28:155-9.
 55. Filatov L, Golubovskaya V, Hurt JC, Byrd LL, Phillips JM, Kaufmann WK. Chromosomal instability is correlated with telomere erosion and inactivation of G2 checkpoint function in human fibroblasts expressing human papillomavirus type 16 E6 oncoprotein. *Oncogene* 1998;16:1825-38.

Non commercial use only